

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation
Non-sterile Solution

Applicator is sterile if package is intact

0.67 ml SEPP® APPLICATOR

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Purposes

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only

Flammable, keep away from fire or flame

- do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch applicator tip
- hold the sponge tip down and pinch the barrel of the applicator only once to activate the ampule and release the antiseptic
- wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.
- do not blot or wipe away
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients USP purified water