

† Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution

Packaging is sterile when in sterile kit

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts	Drug Facts (continued)
<p>Active ingredients</p> <p>Chlorhexidine gluconate 3.15% w/v. Antiseptic Isopropyl alcohol 70% v/v. Antiseptic</p>	<p>Do not use</p> <ul style="list-style-type: none">• on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol• for lumbar puncture or in contact with the meninges• on open skin wounds or as a general skin cleanser• under occlusive patch
<p>Use For the preparation of the patient's skin prior to surgery or injection</p>	<p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>
<p>Warnings</p> <ul style="list-style-type: none">• For external use only• Flammable, keep away from fire or flame• Do not use with electrocautery procedures	

Drug Facts (continued)	Drug Facts (continued)
<p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<ul style="list-style-type: none">• place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute• allow the prepped area to air dry for one and one half minutes (1.5 minutes)• do not blot or wipe dry• discard after a single use
<p>Directions</p> <ul style="list-style-type: none">• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.• maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)• tear open the pouch to expose the swabstick• remove the swabstick from package. Do not touch the foam tip.• prior to surgery or injection, apply swabstick to the procedure site	<p>Other information</p> <ul style="list-style-type: none">• store between 68-77°F (20-25°C)• avoid freezing and excessive heat above 104°F (40°C)• for hospital and professional use only
	<p>Inactive ingredient</p> <p>USP purified water</p>

Each swabstick contains 0.054 fl. oz. (1.6 mL)