

# **+** Prevantics® Swab

REORDER NO. B11400  
NDC 10819-1080-2

[Chlorhexidine Gluconate (3.15%)\*  
and Isopropyl Alcohol (70%) Swab] Antiseptic  
Non-sterile Solution • Packaging is sterile when in sterile kit

\*equivalent to 32 mg of Chlorhexidine Gluconate per pouch

## **Drug Facts**

### **Active ingredients**

Chlorhexidine gluconate 3.15% w/v . . . . .	Antiseptic
Isopropyl alcohol 70% v/v . . . . .	Antiseptic

### **Purpose**

**Use** For the preparation of the patient's skin prior to an injection

### **Warnings**

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

**Allergy alert:** This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash
- If an allergic reaction occurs, stop use and seek medical help right away.

### **Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

**When using this product** keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent



## **Drug Facts** (continued)

injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

### **Other information**

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

**Inactive ingredient** USP purified water

**Questions?** call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

Each Swab contains 0.034 fl. oz. (1 mL)



Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdihc.com



5QB114IS

REORDER NO. S32450

NDC 10819-4077-2

# **†** Prevantics<sup>®</sup> Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

Patient preoperative skin preparation

Non-sterile Solution

Packaging is sterile when in sterile kit

\*equivalent to 50 mg of Chlorhexidine Gluconate per pouch

## **Drug Facts**

### **Active ingredients**

Chlorhexidine gluconate 3.15% w/v. . . . . Antiseptic  
Isopropyl alcohol 70% v/v. . . . . Antiseptic

### **Purpose**

**Use** For the preparation of the patient's skin prior to surgery or injection

### **Warnings**

- For external use only
- Flammable, keep away from fire or flame
- Do not use with electrocautery procedures

## **Drug Facts** (continued)

### **Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:  
• wheezing/difficulty breathing • shock • facial swelling • hives • rash  
If an allergic reaction occurs, stop use and seek medical help right away.

### **Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

## ***Drug Facts*** (continued)

**When using this product** keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### ***Directions***

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply one flat side of foam tip to the proposed skin site and prep the skin in vigorous ▶

## ***Drug Facts*** (continued)

back-and-forth repeated strokes, turning the swabstick over (unused side of the foam tip) halfway in the procedure

- **dry site:** scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds.
- **moist site:** scrub repeatedly for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes.
- do not blot or wipe dry
- discard after a single use

### ***Other information***

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

### ***Inactive ingredient***

USP purified water

### ***Questions?***

call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

**Each swabstick contains 0.054 fl. oz. (1.6 mL)**



Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdihc.com

5RS324IS



REORDER NO. S42850  
NDC 10819-4077-3

# Prevantics<sup>®</sup> Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

Patient preoperative skin preparation

Non-sterile Solution

Packaging is sterile when in sterile kit

\*equivalent to 50 mg of Chlorhexidine Gluconate per pouch

## **Drug Facts**

### **Active ingredients . . . . . Purpose**

Chlorhexidine gluconate 3.15% w/v. . . . . Antiseptic

Isopropyl alcohol 70% v/v. . . . . Antiseptic

**Use** For the preparation of the patient's skin prior to surgery or injection

### **Warnings**

- For external use only
- Flammable, keep away from fire or flame
- Do not use with electrocautery procedures

### **Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include: ▶

## **Drug Facts** (continued)

- wheezing/difficulty breathing • shock • facial swelling • hives • rash
- If an allergic reaction occurs, stop use and seek medical help right away.

### **Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

**When using this product** keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

## **Drug Facts** (continued)

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes, turning the swabstick over (unused side of the foam tip) halfway in the procedure ▶

## **Drug Facts** (continued)

- **dry site:** scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds.
- **moist site:** scrub repeatedly for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes.
- do not blot or wipe dry
- discard after a single use

### **Other information**

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

### **Inactive ingredient**

USP purified water

### **Questions?**

call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

**Each swabstick contains 0.054 fl. oz. (1.6 mL)**



Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 [www.pdihc.com](http://www.pdihc.com)

5RS428IS



# † Prevantics® Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

Patient preoperative skin preparation  
Non-sterile Solution

Packaging is sterile when in sterile kit

\*equivalent to 161 mg of Chlorhexidine Gluconate per pouch

**⚠ WARNING**

**FLAMMABLE**



Keep away from fire or flame.

To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

## Drug Facts

### Active ingredients

Chlorhexidine gluconate 3.15% w/v.....Antiseptic  
Isopropyl alcohol 70% v/v.....Antiseptic

### Purpose

## Drug Facts (continued)

solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)

- do not allow solution to pool
- remove wet materials from prep area

**Use** For the preparation of the patient's skin prior to surgery or injection

### Warnings

For external use only. Flammable, keep away from fire or flame.  
To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until

**Allergy alert:** This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

### Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product

## Drug Facts (continued)

- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

**When using this product** keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

## Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately 8.4 in. x 8.4 in. (21.3 cm by 21.3 cm).
- tear open the pouch to expose the swabstick.
- remove swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes, turning the swabstick over (unused side of the foam tip) halfway in the procedure. ▶

## Drug Facts (continued)

- **dry site:** Scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds.
- **moist site:** Scrub repeatedly for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard after a single use

## Other information

- store between 68 – 77°F (20 – 25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

## Inactive ingredient

USP purified water

## Questions?

call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

**Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)**



Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdihc.com

5SS273IS

